arXiv:math/0307065v1 [math.AG] 4 Jul 2003

[1]

FIBRED KÄHLER AND QUASIPROJECTIVE GROUPS.

FABRIZIO CATANESE
(UNIVERSITÄT BAYREUTH)
THIS ARTICLE IS DEDICATED TO ADRIANO BARLOTTI ON THE OCCASION OF HIS 80-TH BIRTHDAY.

Abstract. We formulate a new theorem giving several necessary and sufficient conditions in order that a surjection of the fundamental group $\pi_1(X)$ of a compact Kähler manifold onto the fundamental group Π_g of a compact Riemann surface of genus $g \geq 2$ be induced by a holomorphic map. For instance, it suffices that the kernel be finitely generated.

We derive as a corollary a restriction for a group G, fitting into an exact sequence $1 \to H \to G \to \Pi_g \to 1$, where H is finitely generated, to be the fundamental group of a compact Kähler manifold.

Thanks to the extension by Bauer and Arapura of the Castelnuovo de Franchis theorem to the quasi-projective case (more generally, to Zariski open sets of compact Kähler manifolds) we first extend the previous result to the non compact case. We are finally able to give a topological characterization of quasi-projective surfaces which are fibred over a (quasi-projective) curve by a proper holomorphic map of maximal rank, and we extend the previous restriction to the monodromy of any fibration onto a curve.

---

[1] The research of the author was performed in the realm of the SCHWERPUNKT "Globale Methode in der komplexen Geometrie", and of the EAGER EEC Project.

Date: February 16, 2003.

1. Introduction

The history of fibrations of algebraic (or Kähler) manifolds $f : X \to C$ over curves C of genus at least 2, called classicaly irrational pencils, has a long history.

Around 1905 almost simultaneously De Franchis and Castelnuovo-Enriques ([D-F05], [Cast05-2]) found that the existence of such a fibration is equivalent to the existence of at least 2 linearly independent holomorphic 1-forms whose wedge product yields a holomorphic 2-form which is identically zero.

Hodge theory was yet to be developed and only much later ([Cat91]) it was shown that combining the Hodge decomposition with the theorem of Castelnuovo De Franchis one obtains a topological characterization of such fibrations via any subspace in De Rham cohomology obtained as the pull-back of a maximal isotropic subspace in the cohomology of C.

Other topological characterizations in terms of the induced surjection of fundamental groups $f_* : \pi_1(X) \to \pi_1(C)$, or other statements in this direction, had been earlier obtained by several authors (cf. Jost-Yau and Siu, [Siu80] , [J-Y83], [J-Y85], [Siu87], who used the theory of harmonic maps, Beauville, [Beau], used instead the generic vanishing theorems of Green and Lazarsfeld, [G-L87]).

In [Cat96] I tried to show how the isotropic subspace theorem, which predicts the genus of the image curve C (it equals to the dimension of the corresponding maximal isotropic subspace), unlike the other statements, could be used to obtain also simple proofs of statements concerning surjections of fundamental groups, as the one given by Gromov ([Gro89]).

Hodge theory also works for quasi-projective manifolds, and it turns out that in the non compact case it works much better than the other methods ([Ba97], [Ara97]). These results were then used in [Cat00] to give topological characterizations of varieties isogenous to a product and of isotrivial fibrations of surfaces.

Kotschick ([Kot99]) instead used very similar methods to give a topological characterization of Kodaira fibrations, which was independently also obtained by Hillman ([Hil00]).

Our first motivation was to extend this result to any fibration, a goal that we have achieved in the following

Theorem 6.4. *Assume that U is a non complete Zariski open set of an algebraic surface and that the following properties hold:*

- *(P1) we have an exact sequence*

$$1 \to \Pi_r \to \pi_1(U) \to \mathbb{F}_g \to 1$$

where $g \geq 2$.

- *(P2) The topological Euler-Poincare' characteristic of U, $e(U)$, equals $2(g-1)(r-1)$.*
- *(P3) For each end $\mathcal{E}$ of U, the corresponding fundamental group $\pi_1^{\mathcal{E}}$ surjects onto a cyclic subgroup of $\mathbb{F}_g$, and each simple geometric generator γ_i has a non trivial image in $\mathbb{F}_g$.*

Then U is a good open set of a fibration, more precisely, there exists a proper holomorphic submersion $f : U \to C$ inducing the previous exact sequence.

To this purpose, we started to put together the existing results, both in the compact and in the non- compact case, with some small addition: we refer to Theorems 4.3 and 5.4 for full statements.

We only indicate here some new results therein:

Theorem A.

Let X be a compact Kähler manifold, and assume that its fundamental group admits a non trivial homomorphism ψ to the fundamental group Π_g of a compact Riemann surface of genus $g \geq 2$, with kernel H.

Then the following conditions are equivalent

- *4) ψ is induced by an irrational pencil of genus g without multiple fibres*
- *5) ψ is surjective and its kernel H is finitely generated*

Theorem A'.

Let X be a compact Kähler manifold and $Y = X - D$ be a Zariski open set. Assume that the fundamental group of Y admits a homomorphism $\psi : \pi_1(Y) \to \mathbb{F}_g$ to a free group of rank g, with kernel H.

Then the following are equivalent

- *1') ψ is induced by a pencil $f : Y \to C$ of type g without multiple fibres and by a surjection $\pi_1(C) \to \mathbb{F}_g$.*
- *2') ψ is surjective and the kernel H of ψ is finitely generated*

The new ingredient here is a remarkable property of free groups and of the fundamental groups of compact curves of genus $g \geq 2$.

This property is that every non trivial **N**ormal subgroup of **I**nfinite index is **N**ot **F**initely generated.

We abbreviate this property by the acronym **NINF**, and we devote section 3 to establishing this result for the above mentioned groups.

This property plays an important role, for instance it shows that, contrary to what is stated by some author, the kernel of the homomorphism between fundamental groups $f_* : \pi_1(X) \to \pi_1(C)$ needs not be finitely generated, it is finitely generated if and only if there are no multiple fibres.

In the case where there are multiple fibres, one can take a ramified base change which eliminates the multiple fibres, and indeed one could extend Theorems A, A' to include the case where H is not finitely generated, but we believed that the results stated in this article are already sufficiently complicated, so we omitted to treat this extension.

We prefered instead to concentrate on some important consequence, concerning the monodromy of fibrations over curves. For instance, putting together Theorem A with the old isotropic subspace theorem we obtain

Corollary 7.3

If a finitely presented group Γ admits a surjection $\Gamma \to \Pi_g$ with finitely generated kernel H, then Γ cannot be the fundamental group of a compact

Kähler manifold X if there is a non zero element $u \in H^1(H,\mathbb{Z})^{\Pi_g}$ such that cup product with u yields the identically zero map

$$H^1(\Pi_g,\mathbb{Z}) \to H^1(\Pi_g, H^1(H,\mathbb{Z})).$$

We then spell out in detail the meaning of degeneracy of the above cup product: it means that there exists a bad monodromy submodule.

By extending everything in the non compact case one obtains then a restriction for the monodromy of fibrations over curves which is in the same spirit as Deligne's semisimplicity Theorem (4.2.6 of [Del 71]).

2. Notation

- Π_g denotes the fundamental group of a compact Riemann surface C_g of genus $g \geq 2$, $\Pi_g := < a_1, \dots a_g, b_1, \dots b_g | [a_1,b_1] \dots [a_g,b_g] = 1 >$.
- $\mathbb{F}_g$ denotes a free group of rank $g \geq 2$
- X will be a compact Kähler manifold

3. Non finitely generated subgroups

This section is devoted to a remarkable property enjoyed, for $g \geq 2$, by the free groups $\mathbb{F}_g$ and by the fundamental groups Π_g of a compact Riemann surface C_g of genus $g \geq 2$.

Definition 3.1. *A group G is said to satisfy property $NINF$ (to be more precise, but less concise, we should call it $NIINFG$) if every* **normal** *non trivial subgroup K of* **infinite index** *is* **non finitely** *generated.*

We shall also say that G is a $NINF$.

The application of the above notion that we shall need is the following

Lemma 3.2. *Let $1 \to A \to \Pi \to B \to 1$ be an exact sequence of group homomorphisms such that*

- *1) A is finitely generated*
- *2) B is infinite*
- *$\phi : \Pi \to B$ factors as $\rho \circ \psi$, where $\psi : \Pi \to G$ is surjective*
- *3) G is a $NINF$*

Then $\rho : G \to B$ is an isomorphism.

Proof. Let $j : A \to \Pi$ be the inclusion and define $K' := ker(\psi \circ j)$, $K := ker(\psi)$. Then $K' = K$ since $K \subset A = ker(\phi)$.

Set $A' := A/K$, so A' injects into $G = \Pi/K$. Moreover, A' is normal in G with quotient $B = \Pi/A$ which is infinite by assumption. Since A' is finitely generated, as a quotient of A, and G is a $NINF$ it follows that A' is trivial.

Whence, $A = K$ and $\rho : G \to B$ is an isomorphism, as desired. □

Lemma 3.3. *A free group $\mathbb{F}_n$ enjoys property $NINF$.*

Proof.

W.l.o.g. assume we may assume $n \geq 2$. We view $\mathbb{F}_n$ as the fundamental group $\pi_1(Y)$, where Y is a bouquet of n circles.

Let Z be the covering space corresponding to a normal subgroup K of infinite index.

Z is indeed the Cayley graph for the infinite group $G := \mathbb{F}_n/K$ with respect to the finite set of n generators, $g_1, \dots g_n$, corresponding to the surjection $\mathbb{F}_n \to G$.

If K is non trivial, then Z is not simply connected, and there is a non trivial minimal closed simplicial path ξ based on the base point $x_0 = 1$: $(x_0 = 1, x_1 = \gamma_1, \dots x_m = \gamma_1 \dots \gamma_m)$, where the γ_i 's belong to the given set of generators $\{g_1, \dots g_n\}$.

Let $M \subset G$ be the set $\{x_0 = 1, x_1, \dots x_m\}$ and let M' be the finite set $MM^{-1} \subset G$.

Since G is infinite , there exist infinitely many h_α's such that $h_\alpha \notin h_\beta M'$ for $\alpha \neq \beta$.

Whence, for $\alpha \neq \beta$ and $\forall x_i, x_j$, we have $h_\alpha x_i \neq h_\beta x_j$. It follows that the cycles $h_\alpha(\xi)$ are homologically independent.

A fortiori, we have shown $Rank(H_1(Z, \mathbb{Z}) = \infty$ and K is not finitely generated.

□

Lemma 3.4. *A fundamental group Π_g enjoys property $NINF$ for $g \geq 2$.*

Proof. Let Γ be a nontrivial normal subgroup of $\Pi := \Pi_g$, and let $f : D \to C$ be the corresponding unramified covering of a compact Riemann surface of genus g. We have $g \geq 2$, whence we may view D as a quotient of the upper half plane $\mathbb{H}$ by the action of the group Γ acting freely and properly discontinuously.

As in [Sieg73], thm. 4, page 35 , we consider fundamental domains, $\mathcal{F}_\Pi$ resp. $\mathcal{F}_\Gamma$, bounded by non-euclidean segments (possibly also lines or half-lines).

While $\mathcal{F}_\Pi$ has finite area, the area of $\mathcal{F}_\Gamma$ is the area of $\mathcal{F}_\Pi$ multiplied by the index of Γ, whence $\mathcal{F}_\Gamma$ has infinite area.

Assume that Γ be finitely generated: then (cf. [Bear83] Thm. 10.1.2 , page 254) there is such a fundamental domain $\mathcal{F}_\Gamma$ with finitely many sides. Since however its area is infinite, it cannot be a non Euclidean ideal polygon, and there are intervals in the real line $\mathbb{P}^1_{\mathbb{R}}$ which need to be added for a compactification of $\mathcal{F}_\Gamma$.

Let us recall the following standard definitions (cf. [Sieg73], and especially [Se-So92], [Bear83])

Definition 3.5.
- *1) A subgroup Γ of $PSL(2, \mathbb{R})$ acting properly discontinuosly on $\mathbb{H}$ is called a Fuchsian group (more generally, a Fuchsian group is a conjugate in $PSL(2, \mathbb{C})$ of such a subgroup).*
- *2) Γ is properly discontinuos if and only if it is discrete in $PSL(2, \mathbb{R})$*
- *3) The* **Limit Set** *$L(\Gamma)$ is defined as*

$$L(\Gamma) := \cup_{z \in \mathbb{H}} (\overline{\Gamma z} \cap \mathbb{P}^1_{\mathbb{R}}).$$

- *4) Equivalently, (cf.* [Se-So92]*, page 108)*

$$L(\Gamma) := \{z \in \mathbb{P}^1_{\mathbb{R}} | \exists \gamma \in \Gamma, \gamma \neq 1, s.t. \gamma(z) = z\}.$$

- *5) A Fuchsian subgroup Γ is said to be of the I Kind if $L(\Gamma) = \mathbb{P}^1_{\mathbb{R}}$ (else, it is said to be of the II Kind).*

We have (cf. e.g. Lemma 3.12.2, page 108 of [Se-So92]) that if $\mathbb{H}/\Gamma$ is compact, then Γ is of the I Kind. This implies a consequence for our group Π.

In fact, since Π normalizes Γ, then Π carries the limit set $L(\Gamma)$ to itself.

Let I be an interval in the real line $\mathbb{P}^1_{\mathbb{R}}$ which is in the boundary of $\mathcal{F}_{\Gamma}$. Since Π is of the I Kind, there is $x \in I$ and $g \in \Pi - \{1\}$ such that $gx = x$.

Since, as we observed, $g(L(\Gamma)) = L(\Gamma)$, g carries the interior of the complement to $L(\Gamma)$ in $\mathbb{P}^1_{\mathbb{R}}$ to itself. Let the interval (a, b) be the connected component of this interior containing x. Since moreover $gx = x$, g carries (a, b) to itself.

Assume that $a \neq b$: then g^2 has three fixed points (a, b, x), thus g^2 is the identity, contradicting the hyperbolicity of g.

If however $a = b$, this means that the limit set $L(\Gamma)$ consists of a single point a (fixed by each $g \in \Pi$) : contradicting that Γ is of the I Kind.

Q.E.D.

4. Mappings to curves

If a manifold X is fibred (with connected fibres) onto a curve C, certainly we have a surjection of fundamental groups $\pi_1(X) \to \pi_1(C)$.

In fact, let $p_1, \dots p_r$ be the critical values of f, and set $C^* := C - \{p_1, \dots p_r\}$, $X^* := f^{-1}(C^*)$: then we have surjections $\pi_1(X^*) \to \pi_1(X)$, $\pi_1(C^*) \to \pi_1(C)$, and an exact homotopy sequence

$$\pi_1(F) \to \pi_1(X^*) \to \pi_1(C^*) \to 1.$$

It suffices to observe that the surjection $\pi_1(X^*) \to \pi_1(C)$ factors through $\pi_1(X) \to \pi_1(C)$.

If however there is a non trivial holomorphic map $F : C \to C'$, such that F does not factor as $F = F' \circ F"$, where F' is unramified, then it is not difficult to show that there is a surjection of fundamental groups $F_* : \pi_1(C) \to \pi_1(C')$.

This is the reason why one needs some extra assumptions on a surjection of fundamental groups $\pi_1(X) \to \pi_1(C')$ in order to decide whether the corresponding map is a fibration (i.e., it has connected fibres).

Recall (cf. e.g. [C-K-O-03], Lemma 3 , page 283) the following

Definition 4.1. *Let m_i ,$\forall i = 1, \dots r$, be the G.C.D. of the multiplicities of the components of the divisor $f^{-1}(p_i)$ ($p_1, \dots p_r$ are again the critical values of f).*

Then the orbifold fundamental group $\pi_1^{orb}(f)$ is defined as the quotient of $\pi_1(C - \{p_1, \dots p_r\})$ by the subgroup normally generated by $\{\gamma_i^{m_i}\}$, γ_i being a simple geometric path around the point p_i.

As a corollary of the results of the previous section we have

Lemma 4.2. *If X admits a surjective holomorphic map f with connected fibres $f : X \to C$ where C is a Riemann surface of genus $g \geq 0$, then the induced homomorphism $f_* : \pi_1(X) \to \Pi_g$ is surjective, and its kernel H is finitely generated exactly when $g = 0$ or when $g \geq 1$ and there are no multiple fibres, i.e., $\pi_1^{orb}(f) \cong \pi_1(C)$.*

Proof. As well known (cf. e.g. a slightly general version given [C-K-O-03], Lemma 3 , page 283, whose notation we will follow) we have an exact sequence

$$\pi_1(F) \to \pi_1(X) \to \pi_1^{orb}(f) \to 1$$

where F is a smooth fibre of f, and, $p_1, \dots p_r$ being the critical values of f, and $m_i, \forall i = 1, \dots r$, being the respective G.C.D. of the multiplicities of the components of the divisor $f^{-1}(p_i)$,the orbifold fundamental group $\pi_1^{orb}(f)$ is defined as the quotient of $\pi_1(C - \{p_1, \dots p_r\})$ by the subgroup normally generated by $\{\gamma_i^{m_i}\}$, γ_i being a simple geometric path around the point p_i.

Let $\psi := f_*$. Thus $ker(\psi)$ contains the normal subgroup K, image of $\pi_1(F)$, which is finitely generated since F is compact, and the cokernel $ker(\psi)/K$ is isomorphic to the kernel of $\rho : \pi_1^{orb}(f) \to \pi_1(C)$.

Therefore $ker(\psi)$ is finitely generated if and only if $ker\rho$ is finitely generated. This is then the case for $g = 0$, so let us assume that $g \geq 1$. If we moreover assume that there are no multiple fibres, then ρ is an isomorphism, and we are again done.

Otherwise, $\pi_1^{orb}(f)$ is a Fuchsian group and the same proof as in Lemma 3.4 shows that $\pi_1^{orb}(f)$ is $NINF$. Then $ker\rho$ is finitely generated if only if it is trivial: since the alternative that its index be finite is ruled out by the condition $g \geq 1$.

Finally, if $ker\rho$ is trivial, then ρ is an isomorphism, and by looking at the Abelianization we see that $r = 1$. But then the orbifold fundamental group $\pi_1^{orb}(f)$ is a free product of a free group of rank $2g - 1$ with a cyclic group of order m_1 and its Abelianization is then not a free Abelian group of rank $2g$, a contradiction.

Q.E.D.

Theorem 4.3. *Let X be a compact Kähler manifold, and assume that its fundamental group admits a non trivial homomorphism ψ to the fundamental group Π_g of a compact Riemann surface of genus $g \geq 2$, with kernel H.*

Then the following conditions are equivalent

- *1) ψ is induced by an irrational pencil of genus g i.e. there is a surjective holomorphic map f with connected fibres $f : X \to C$ such that $\psi = f_*$, and where C is a Riemann surface of genus g.*
- *2) ψ is surjective and the image of $\psi^* : H^1(\Pi_g, \mathbb{Q}) \to H^1(X, \mathbb{Q})$ contains a g-dimensional maximal isotropic subspace (for the bilinear pairing $H^1(X, \mathbb{Q}) \times H^1(X, \mathbb{Q}) \to H^2(X, \mathbb{Q})$)*
- *3) ψ induces an injective map in cohomology $\psi^* : H^1(\Pi_g, \mathbb{Q}) \to H^1(X, \mathbb{Q})$, and the image of ψ^* contains a g-dimensional maximal isotropic subspace*

likewise, the following conditions are also equivalent to each other

- *4) ψ is induced by an irrational pencil of genus g without multiple fibres, i.e. , for each fibre F' the equation of divisors $F' = rD$, with $r \geq 1$, implies $r = 1$.*
- *5) ψ is surjective and its kernel H is finitely generated*

Proof. We observe first that 2) implies 3) follows since a surjective homomorphism induces a surjective homomorphism between the abelianizations, and dualizing one obtains an injective homomorphism in cohomology.

Recall that, by the isotropic subspace theorem of [Cat91], given a maximal isotropic subspace $V \subset H^1(X, \mathbb{Q})$, of dimension g, there is a holomorphic fibration onto a curve C of genus g such that $V \subset f^*(H^1(C, \mathbb{Q}))$.

Now, 1) implies 2) because, if the pull back of a maximal isotropic subspace from the given curve C is not maximal, then we have another fibration $f' : X \to C'$ to a curve of genus $g' > g$ such that $f^*(H^0(C, \Omega^1_{C'})) \subset f^*(H^0(C', \Omega^1_{C'})$, whence f factors through f', contradicting the fact that f has connected fibres.

Let us show that 3) implies 1). The isotropic subspace theorem gives us the desired $f : X \to C$, where C has genus g.

Since however C is a classifying space for Π_g, there is a continuous map $F : X \to C$ such that $\psi = F_*$.

Compose both maps with the Jacobian embedding $\alpha : C \to J$ and observe that by the proof of the isotropic subspace theorem the two subspaces $f^*(H^1(C, \mathbb{Q}))$ and $F^*(H^1(C, \mathbb{Q}))$ coincide.

Therefore the two maps $\alpha \circ F$,$\alpha \circ f$ are given by integrals of the same differentiable 1-forms, hence there is an isogeny $p : J \to J$ such that $\alpha \circ f = p \circ \alpha \circ F$. We get thus, up to changing F in its homotopy equivalence, a factorization $f = p' \circ F$. Thus, p' is surjective, and actually it has degree 1 or otherwise $f^*(\eta)$, with (η) the positive generator of $H^2(C, \mathbb{Z})$), would be divisible, contradicting that f has connected fibres.

The conclusion is that f and F are homotopy equivalent, thus $\psi = F_* = f_*$.

The implication 4) → 5) is exactly lemma 4.2, so we are left with showing that 5) implies 4).

Now, 5) implies that the image of ψ^* contains a g-dimensional isotropic subspace. Assume this subspace not to be maximal: then there is a fibration $f : X \to C$ where the genus g' of C is strivtly larger than g.

Argueing as we did before, we find a factorization of ψ through f'_*.

Since ψ is surjective, Lemma 3.2. applies and we get that $g' = g$, and $\psi = f_*$.

Finally, f has no multiple fibres again by Lemma 4.2.

Q.E.D.

5. THE LOGARITHMIC CASE

In this section we shall generalize the results of the previous section to the case where we have a Zariski open set Y in a compact Kähler manifold X. One may assume w.l.o.g. that the complement $X - Y$ is a normal crossings divisor D. We shall consider holomorphic maps $f : Y \to C$, where C is Zariski open in a compact curve $\bar{C}$, and the map f is meromorphic on X, whence there is another compactification $\bar{X}$ of X where f extends holomorphically.

When we shall say that f is a **pencil**, we shall mean that f is as above, that the extension $\bar{f}$ of f has connected fibres, and that f is surjective (Arapura calls these maps **admissible maps**). We shall denote by B the complement $\bar{C}-C$, because quite often it will be the branch locus of a fibration of a compact manifold.

However, X will not necessarily be non compact, the reason for this being that we shall here consider surjective homomorphisms $\pi_1(X) \to \mathbb{F}_n$ to a non Abelian free group.

Notice moreover that

- any automorphism of Π_g composed with the standard surjection $p : \Pi_g \to \mathbb{F}_g$ such that $p(a_i) = p(b_i) = x_i$ produces a maximal isotropic subspace of $H^1(C, \mathbb{Z})$.
- there is a surjection $p : \Pi_g \to \mathbb{F}_n$ iff $g \geq n$ (since $Im(p^*)$ is an isotropic subspace of dimension n).

The next theorem extends the results of I. Bauer and D. Arapura (Thms 2.1. and 3.1 of [Ba97] and Cor. 1.8 of [Ara97], cf. also thm. 2.11 of [Cat00]) using the new ideas introduced in the previous sections.

Observe that also in this context a pencil induces a surjective homomorphism of fundamental groups.

Definition 5.1. *Let $f : Y \to C$ be a pencil as above. We shall say that f is of type g if either*

- *C is compact of genus g (then $\pi_1(C) \cong \Pi_g$)*
- *C is not compact and its first Betti number equals g (then $\pi_1(C) \cong \mathbb{F}_g$).*

Definition 5.2. *Let $f : Y \to C$ be a pencil as above.*

We may assume that f extends to a holomorphic fibration $F : X \to \bar{C}$. We can separate the complementary divisor $D = X - Y$ into three parts :

- *D^{hor} : the union of the components dominating C*
- *D^{FVert}: the union of the fibres over $\bar{C} - C$*
- *D^{PVert}: the union of the components mapping to points of C.*

We define then the orbifold fundamental group of f by the usual procedure: let $p_1, \dots p_r$ be the critical values of f, and $m_i, \forall i = 1, \dots r$, be the respective G.C.D. of the multiplicities of the components of the divisor $f^{-1}(p_i)$. Then the orbifold fundamental group $\pi_1^{orb}(f)$ is defined as the quotient of $\pi_1(C - \{p_1, \dots p_r\})$ by the subgroup normally generated by $\{\gamma_i^{m_i}\}$, γ_i being a simple geometric path around the point p_i.

Lemma 5.3. *Let X be a compact Kähler manifold and $Y = X - D$ be a Zariski open set. If Y admits a pencil $f : Y \to C$, then the induced homomorphism $f_* : \pi_1(X) \to \pi_1(C)$ is surjective, and its kernel H is finitely generated exactly when $g = 0$ or when $g \geq 1$ and there are no multiple fibres, i.e., $\pi_1^{orb}(f) \cong \pi_1(C)$.*

Proof. We use once more the exact sequence (the situation being more general, but the proof exactly the same as in [C-K-O-03], Lemma 3 , page 283)

$$\pi_1(F) \to \pi_1(X) \to \pi_1^{orb}(f) \to 1$$

where F is a smooth fibre of f which is transversal to D^{hor}.

Let $\psi := f_*$. Thus $ker(\psi)$ contains the normal subgroup K, image of $\pi_1(F)$, which is finitely generated since F is of finite type, and the cokernel $ker(\psi)/K$ is isomorphic to the kernel of $\rho : \pi_1^{orb}(f) \to \pi_1(C)$.

Therefore $ker(\psi)$ is finitely generated if and only if $ker\rho$ is finitely generated. Assume that there are no multiple fibres: then ρ is an isomorphism.

Otherwise, $\pi_1^{orb}(f)$ is a Fuchsian group and the same proof as in Lemma 3.4 shows that $\pi_1^{orb}(f)$ is $NINF$. Then $ker\rho$ is finitely generated if only if it is trivial, its index being infinite for $g \geq 1$.

Q.E.D.

Theorem 5.4. *Let X be a compact Kähler manifold and $Y = X - D$ be a Zariski open set. Assume that the fundamental group of Y admits a homomorphism $\psi : \pi_1(Y) \to \mathbb{F}_g$ to a free group of rank g, with kernel H.*

Then the following are equivalent

- *1) ψ is induced by a pencil $f : Y \to C$ of type g and by a surjection $\pi_1(C) \to \mathbb{F}_g$.*
- *2) ψ is surjective and the image of $\psi^* : H^1(\mathbb{F}_g, \mathbb{Q}) \to H^1(Y, \mathbb{Q})$ is a g-dimensional maximal isotropic subspace (for the bilinear pairing $H^1(Y, \mathbb{Q}) \times H^1(Y, \mathbb{Q}) \to H^2(Y, \mathbb{Q})$)*

likewise, the following are also equivalent to each other

- *1') ψ is induced by a pencil of type g without multiple fibres*
- *2') ψ is surjective and the kernel H of ψ is finitely generated*

Finally, the curve C is compact if and only if $\psi^*(H^1(\mathbb{F}_g, \mathbb{Q}))$ is also an isotropic subspace in $H^1(X, \mathbb{Q})$ $(\subset H^1(Y, \mathbb{Q}))$.

Proof.

1) implies 2) : it suffices to show that $\psi^*(H^1(\mathbb{F}_g, \mathbb{Q}))$ is a maximal isotropic subspace. Assume the contrary : then, there is a strictly larger maximal isotropic subspace V induced (cf.[Cat00], thm. 2.11) by a pencil $f' : Y \to C'$.

The pencil is induced by integrations of linearly independent forms in $H^0(\Omega^1(logD)$, whence we get a factorization $Y \to C' \to C$: since f has connected fibres $C' \cong C$, contradicting that the logarithmic genus of C' is strictly larger than g.

Assume 2): then by Arapura's cor. 1.9 there is a pencil $f : Y \to C$ and $\tau : H_1(C, \mathbb{Z}) \to \mathbb{Z}^g$ such that there is a factorization in homology $H_1(\psi) = \tau \circ H_1(f)$.

Since ψ is surjective, then also $H_1(\psi)$ is such, whence we may lift τ to a surjection $\pi_1(C) \to \mathbb{F}^g$ (since $Aut(\mathbb{F}_g)$ surjects onto $GL(g, \mathbb{Z})$, cf. [MKS66], sections 3.5 and 3.6).

The equivalence of 1') and 2') follows as in the proof of Theorem 4.3 in view of Lemma 5.3 . The last assertion is already contained in theorem. 2.11, loc. cit.

Q.E.D.

6. Fibred algebraic surfaces and good open sets

In this section we shall consider a smooth compact algebraic surface, and a holomorphic fibration $f : S \to C$.

Definition 6.1. *A* **good open set** *of a fibration will be any set of the form:* $U = f^{-1}(C - B)$, *where* B *is any finite set containing the set of critical values of* f.

Remark 6.2. *In the above situation one has an exact sequence of fundamental groups*

$$1 \to \pi_1(F) \to \pi_1(U) \to \pi_1(C - B) \to 1$$

where F *is any fibre of* f *over a point of* $C - B$.

The next theorem will give a topological characterization of good open sets of some fibration.

The case where $U = S$ was already treated by Kotschick ([Kot99], Proposition 1) and Hillman ([Hil00]), cf. also Kapovich ([Kap98]), and we only prove an ϵ^2 more general result:

Theorem 6.3. *Assume that* S *is a compact Kähler surface and that we have an exact sequence*

$$1 \to \Pi_r \to \pi_1(S) \to \Pi_g \to 1$$

where $g \geq 2$. *If moreover the topological Euler-Poincare' characteristic of* S, $e(S)$, *equals* $4(g-1)(r-1)$, *then there exists a holomorphic submersion* $f : S \to C$ *inducing the previous exact sequence.*

Proof. By Theorem 4.3 we find a fibration $f : S \to C$, where C has genus g, inducing the given epimorphism of fundamental groups.

By the theorem of Zeuthen-Segre the Euler-Poincare' characteristic of S, $e(S)$, equals $4(g-1)(s-1)+\mu$ where s is the genus of a smooth fibre of f and where $\mu \geq 0$, equality holding if and only if all the singular fibres are multiple of a smooth elliptic curve. It is clear that $s \geq r$.

Our assumption implies $4(g-1)(s-r)+\mu = 0$, whence $s = r$ and $\mu = 0$. We are therefore done in the case where $r = 0$ or $r \geq 2$.

If finally $r = 1$, we are done unless there is some multiple fibre. But the existence of a multiple fibre is excluded by Lemma 4.2.

Q.E.D.

We come now to the non compact case

Theorem 6.4. *Assume that* U *is a non complete Zariski open set of an algebraic surface and that the following properties hold:*

- *(P1) we have an exact sequence*
  $$1 \to \Pi_r \to \pi_1(U) \to \mathbb{F}_g \to 1$$
  where $g \geq 2$.
- *(P2) The topological Euler-Poincare' characteristic of* U, $e(U)$, *equals* $2(g-1)(r-1)$.
- *(P3) For each end* $\mathcal{E}$ *of* U, *the corresponding fundamental group* $\pi_1^{\mathcal{E}}$ *surjects onto a cyclic subgroup of* $\mathbb{F}_g$, *and each simple geometric generator* γ_i *has a non trivial image in* $\mathbb{F}_g$.

Then U *is a good open set of a fibration, more precisely, there exists a proper holomorphic submersion* $f : U \to C$ *inducing the previous exact sequence.*

Proof. By Thm. 5.4 we have a fibration $f : U \to C$ inducing the surjection $\pi_1(U) \to \mathbb{F}_g \to 1$, and w.l.o.g. we have an extension $\bar{f} : \bar{S} \to \bar{C}$, where $\bar{S}$ is a blow-up of S. By condition (P3) there is no component of D which is horizontal, and each component of D pulls back to a fibre of $\bar{f}$.

It turns therefore out that there is no point of indeterminacy of f on D, whence $\bar{S} = S$, and again by condition (P3) U is the full inverse image of C under $\bar{f}$.

It suffices to apply the logarithmic version of the Zeuthen Segre theorem, similarly to thm. 2.14 of [Cat00], and we conclude that ($e(U)$ being the same in ordinary and Borel-Moore homology by virtue of Poincare' duality) $e(U) = 2(g-1)(r-1) \geq 2(g-1)(s-1) + \mu$, where $s \geq r$ is the genus of a smooth fibre of f. Whence, as usual, $0 \geq 2(g-1)(s-r) + \mu$, thus $s = r$ and $\mu = 0$. We conclude as in theorem 6.3.

Q.E.D.

Note. The next question : when is the fibration f a constant moduli fibration? was already answered, with similar methods, in the previous paper [Cat00], cf. 5.4 and 5.7.

7. Restrictions for the monodromy

Before we present some interesting corollary of the previous theorem, we need to recall some well known results

Lemma 7.1. *Let* X *be a topological manifold and* Γ *its fundamental group. Then* $H^1(X, \mathbb{Z}) = H^1(\Gamma, \mathbb{Z})$ *and* $H^2(\Gamma, \mathbb{Z})$ *injects into* $H^2(X, \mathbb{Z})$.

Proof. Let $\tilde{X}$ be the universal covering of X, so that $X \cong \tilde{X}/\Gamma$.

The proof is a direct consequence of the spectral sequence for group cohomology with terms $H^p(\Gamma, H^p(\tilde{X}, \mathbb{Z}))$, converging to a suitable graded quotient of $H^{p+q}(X, \mathbb{Z})$, in view of the fact that $H^1(\tilde{X}, \mathbb{Z}) = 0$.

□

Lemma 7.2. *Let* $1 \to H \to \Gamma \to B \to 1$ *be an exact sequence of groups, where* B *is a finitely generated free group, or the fundamental group* Π_g *of a compact hyperbolic Riemann surface. Assume that:*

(**) $H^2(B, \mathbb{Z})$ *injects into* $H^2(\Gamma, \mathbb{Z})$.

Then $H^1(B,\mathbb{Z}) \subset H^1(\Gamma,\mathbb{Z})$ *with quotient* $H^1(H,\mathbb{Z})^B$*, and the cup product* $H^1(B,\mathbb{Z}) \times H^1(\Gamma,\mathbb{Z}) \to H^2(\Gamma,\mathbb{Z})$ *lands into the subgroup* F *fitting into the exact sequence* $0 \to H^2(B,\mathbb{Z}) \to F \to H^1(B, H^1(H,\mathbb{Z})) \to 0$.

In particular, let V *a maximal isotropic subspace of* $H^1(B,\mathbb{Z})$*: then* V *remains a maximal isotropic subspace in* $H^1(\Gamma,\mathbb{Z})$

only if (resp. : if and only if, in the case where B *is free)*

(***) the cup product $H^1(B,\mathbb{Z}) \times H^1(H,\mathbb{Z})^B \to H^1(B, H^1(H,\mathbb{Z}))$ *is non degenerate in the second factor.*

Proof.

The proof of the first assertions is a direct consequence of the spectral sequence for group cohomology, with terms $H^p(B, H^q(H,\mathbb{Z}))$, converging to a suitable graded quotient of $H^{p+q}(\Gamma,\mathbb{Z})$, in view of the fact that , by assumption (**), the differential $d_2 : H^1(H,\mathbb{Z})^B = H^0(B, H^1(H,\mathbb{Z})) \to H^2(B,\mathbb{Z})$ is zero.

The second assertion holds with "if and only if" in the case where B is a free group, since then $H^2(B,\mathbb{Z}) = 0$, $F = H^1(B, H^1(H,\mathbb{Z}))$, and the question is whether $H^1(B,\mathbb{Z})$ is a maximal isotropic subspace in $H^1(\Gamma,\mathbb{Z})$.

In the other case, observe that $H^2(B,\mathbb{Z}) = \mathbb{Z}$, and that we can find two maximal isotropic subspaces V, V' such that $H^1(B,\mathbb{Z}) = V \bigoplus V'$: moreover then the cup product yields an isomorphism of V' with $V^\vee$.

If we get an element $w' \in H^1(H,\mathbb{Z})^B$ annihilating $H^1(B,\mathbb{Z})$, this means that there is a lift $w \in H^1(\Gamma,\mathbb{Z}) - H^1(B,\mathbb{Z})$ such that $w \cup H^1(B,\mathbb{Z}) \subset H^2(B,\mathbb{Z})$. In particular, there is $u \in V^\vee$ such that $(w-u) \cup V = 0$.

We easily conclude then that the span of V and of $w-u$ is isotropic.

□

Corollary 7.3. *If the finitely presented group* Γ *admits a surjection* $\Gamma \to \Pi_g$ *with finitely generated kernel* H*, then* Γ *cannot be the fundamental group of a compact Kähler manifold* X *if there is a non zero element* $u \in H^1(H,\mathbb{Z})^{\Pi_g}$ *such that cup product with* u *yields the identically zero map*

$$H^1(\Pi_g,\mathbb{Z}) \to H^1(\Pi_g, H^1(H,\mathbb{Z})).$$

We want now to explicitly write down, for Π equal either to Π_g or to a free group $\mathbb{F}_g$, the condition that there is a non zero element $u \in H^1(H,\mathbb{Z})^\Pi$ such that cup product with u yields the identically zero map

$$H^1(\Pi,\mathbb{Z}) \to H^1(\Pi, H^1(H,\mathbb{Z})).$$

Observe first that $H^1(\Pi,\mathbb{Z}) = Hom_{\mathbb{Z}}(\Pi,\mathbb{Z}) \cong \mathbb{Z}^b$, where $b = g$ in the free case, otherwise $b = 2g$.

The condition that, $\forall \phi \in Hom_{\mathbb{Z}}(\Pi,\mathbb{Z})$, $\phi u = 0$ in $H^1(\Pi, H^1(H,\mathbb{Z}))$ means that there is an element $v_\phi \in H^1(H,\mathbb{Z}))$ such that

$$\forall \gamma \in \Pi,\ \phi(\gamma) u = \gamma v_\phi - v_\phi.$$

Taking a basis $\phi_1, \dots \phi_b$, we get $v_1, \dots v_b$ such that

$$(1)\ \forall \gamma \in \Pi,\ \gamma v_j = v_j + \phi_j(\gamma) u.$$

Recall moreover that u is invariant, whence

$$(2)\ \forall \gamma \in \Pi,\ \gamma u = u.$$

Conditions (1), (2), and the $\mathbb{Z}$-linear independence of the characters ϕ_j implies the $\mathbb{Z}$-linear independence of $u, v_1, \dots v_b$ since the $\mathbb{Z}$-module $H^1(H, \mathbb{Z}))$ is torsion free.

Definition 7.4. *A bad monodromy module is a free $\mathbb{Z}$-module of rank $b+1$, with basis $u, v_1, \dots v_b$, and with an action of Π given by (1) and (2).*

Example 7.5. *Let H be a finitely generated group, and let $1 \to H \to \Gamma \to \Pi_g \to 1$ be an exact sequence such that the induced action of Π_g on H by conjugation induces on the $\mathbb{Z}$ dual of the Abelianization of H a Π_g -module structure which contains a bad monodromy module. Then Γ cannot be the fundamental group of a compact Kähler manifold.*

Remark 7.6. *One can use the same type of restriction in the case where $U \neq X$ is the inverse image of the non critical values of a fibration $f : X \to \bar{C}$, and obtain in this way a restriction for the monodromy in the case where $\bar{C} - B$ has first Betti number at least 2.*

Remark 7.7. *To see finally the relation of the above condition with the theory of Lefschetz pencils (in particular with the splitting in invariant and vanishing cycles), let us observe that our cup product is non degenerate if we have a monodromy invariant splitting $H^1(H, \mathbb{Z}) = H^1(H, \mathbb{Z})^{\Pi} \bigoplus W$.*

Because then we may write $v_\phi = u_\phi + w_\phi$ and we obtain

$$\forall \gamma \in \Pi,\ \phi(\gamma) u = \gamma v_\phi - v_\phi = \gamma w_\phi - w_\phi \in W,$$

whence $u = 0$.

This splitting is proven by Deligne's Semi-simplicity Theorem (4.2.6 of [Del 71]*, cf. also thm. 9.1, page 37 , Chapter II of* [Grif 84]*).*

References

[ABCKT96] J. Amoros, M. Burger , K. Corlette, D. Kotschick, D. Toledo "Fundamental groups of compact Kähler manifolds ", Amer. Math. Soc. Mathematical Surveys and Monographs , vol. 44 (1996).

[Ara97] D. Arapura , " Geometry of cohomology support loci for local systems I. " , J. Alg. Geom. 6,3 (1997), 563-597.

[B-P-V84] W. Barth - C. Peters - A. van de Ven, "Compact complex surfaces", Springer Ergebnisse F.3, vol. 4 (1984), Berlin Heidelberg

[Ba97] I. Bauer, "Irrational pencils on non-compact algebraic manifolds", Int. Jour. of Math. 8, 4 (1997), 441-450.

[Bear83] A.F. Beardon, "The geometry of discrete groups", Springer Verlag GTM (1983),

[Beau] A. Beauville, Appendix to [Cat91], Inv. Math. 104, 289 (1991).

[Cast05-2] G. Castelnuovo, " Sulle superficie aventi il genere aritmetico negativo", Rendiconti del Circ. Mat. di Palermo, t. XX (1905), pages 501-506 of the Memorie Scelte.

[Cast] G. Castelnuovo, " Memorie scelte", Zanichelli, Bologna (1937).

[Cat91] F. Catanese, "Moduli and classification of irregular Kähler manifolds (and algebraic varieties) with Albanese general type fibrations", Inv. Math. 104, 263-289 (1991), with an appendix by A. Beauville, page 289.

[Cat96] F. Catanese, "Fundamental groups with few relations", in 'Higher dimensional complex varieties", Proc. Trento 1994, Andreatta and Peternell editors, de Gruyter, Berlin (1996), 163-165.

[Cat00] F. Catanese, " Fibred surfaces, varieties isogenous to a product and related moduli spaces", Amer. J. Math. 122 (2000), 1-44.

[C-K-O-03] F. Catanese, JH Keum, K. Oguiso " Some remarks on the universal cover of an open K3 surface", Math.Ann. 325 (2003), 279-286.

[Com22] A. Comessatti, "Intorno alle superficie algebriche irregolari con $p_g \geq 2(p_a+2)$ e ad un problema analitico ad esse collegato", Rend. Circ. Mat. di Palermo, 46 (1922), 1 - 48.

[D-F05] M. de Franchis, " Sulle superficie algebriche le quali contengono un fascio irrazionale di curve ", Rendiconti Circ. Mat. di Palermo, XX (1905), 49-54.

[Del 71] P. Deligne, " Theprie de Hodge : II ", Publ. Math. I.H.E.S., 40, 2 (1971), 5-57.

[E-S64] J. Eells - J. Sampson, "Harmonic mappings of Riemannian manifolds", Amer. J. Math. 85 (1964), 109-160.

[Fen48] W. Fenchel, "Estensioni di gruppi discontinui e trasformazioni periodiche delle superficie", Atti Acc. Naz. Lincei, Rend. Sc. fis. mat. e nat. serie 8, vol V (1948), 326-329.

[G-L87] M. Green - R. Lazarsfeld, "Deformation theory, generic vanishing theorems, and some cojectures of Enriques, Catanese and Beauville", Inv. Math. 90 (1987), 389-407.

[G-L91] M. Green - R. Lazarsfeld, "Higher obstructions to deforming cohomology groups of line bundles", J. Amer. Soc. 4 (1991), 87-103.

[Grif 84] P. Griffiths, "Topics in transcendental Algebraic Geometry", Annals of Math. Studies 106, (1984), Princeton Univ. Press.

[Gro89] M. Gromov, "Sur le groupe fondamental d'une varieté Kählerienne, C. R. Acad. Sci. Paris 308, DI (1989), 67- 70.

[Gro91] M. Gromov, "Kähler hyperbolicity and L^2-Hodge theory", J. Diff. Geom. 33 (1991), 263-292.

[Hil00] J.A. Hillman, " Complex surfaces which are fibre bundles", Topology Appl. 100 (2000), no. 2-3, 187–191

[J-Y83] J. Jost, S.T. Yau "Harmonic mappings and Kähler manifolds", Math. Ann. 262 (1983), 145-166.

[J-Y85] J. Jost, S.T. Yau "A strong rigidity theorem for a certain class of compact complex analytic surfaces", Math. Ann. 271 (1985), 143-152.

[J-Y93] J. Jost, S.T. Yau "Applications of quasilinear P.D.E.'s to algebraic geometry and arithmetic lattices ", in 'Algebraic Geometry and related topics', Proc. Symp. Inchoen 1992, Yang et al ed. (1993), Int. Press Boston, 169-190.

[Kap98] M. Kapovich, " On normal subgroups in the fundamental groups of complex surfaces", GT/9808085.

[Ker83] S. Kerchoff, "The Nielsen realization theorem", Bull. AMS 2 (1980), 452-454, Annals of Math. 117 (1983), 235-265.

[Kot99] D. Kotschick, " On regularly fibered complex surfaces.", Proceedings of the Kirbyfest (Berkeley, CA, 1998), 291–298 (electronic), Geom. Topol. Monogr., 2, Geom. Topol. Publ., Coventry, (1999).

[MKS66] W. Magnus, A. Karrass, D. Solitar "Combinatorial group theory", Interscience Publishers (1966) (Dover reprint 1976).

[Mok85] N. Mok, "The holomorphic or anti-holomorphic character of harmonic maps into irreducible compact quotients of polydiscs", Math. Ann. 272 (1985), 197-216.

[Mok88] N. Mok, "Strong rigidity of irreducible quotients of polydiscs of finite volume", Math. Ann. 282 (1988), 555-578.

[Mos73] G. Mostow, "Strong rigidity of locally symmetric spaces", Ann. Math. Studies 78, Princeton Univ. Press 1973.

[Se-So92] M. Seppala, T. Sorvali, "Geometry of Riemann surfaces and Teichmller spaces", North-Holland Mathematics Studies, 169. North-Holland Publishing Co., Amsterdam, (1992) ii+263 pp.

[Sieg73] C.L. Siegel, "*Topics in complex function theory, Vol. II*' Interscience Tracts in pure and appl. Math., n. 25, Wiley, N.Y. (1973).

[Siu80] Y.T. Siu, "The complex analyticity of harmonic maps and the strong rigidity of compact Kähler manifolds", Annals of Math. 112 (1980), 73-111.

[Siu87] Y.T. Siu, "Strong rigidity for Kähler manifolds and the construction of bounded holomorphic functions", in 'Discrete groups in geometry and analysis', R. Howe ed., Birkhauser 1987, 124-151.

[Tro96] A. Tromba, "Dirichlet's energy on Teichmüller's moduli space and the Nielsen realization problem", Math. Z. 222 (1996), 451-464.

Author's address:

Prof. Fabrizio Catanese
Lehrstuhl Mathematik VIII
Universität Bayreuth, NWII
D-95440 Bayreuth, Germany

e-mail: Fabrizio.Catanese@uni-bayreuth.de